



This is verified as Dr. Jean Dodds' Vaccination Protocol:

CANINE VACCINATION PROTOCOL – 2009

MINIMAL VACCINE USE

W. Jean Dodds, DVM
HEMOPET
938 Stanford Street
Santa Monica, CA 90403

 310-828-4804 

*Fax: 310/ 453-5240 ** Please Note -- NEW FAX # for Home Office ***

e-mail: hemopet at hotmail com
(remove spaces, use @ symbol and .com)

Note: The following vaccine protocol is offered for those dogs where minimal vaccinations are advisable or desirable. The schedule is one I recommend and should not interpreted to mean that other protocols recommended by a veterinarian would be less satisfactory. It's a matter of professional judgment and choice.

Age of Pups	Vaccine Type
9 - 10 weeks	Distemper + Parvovirus, MLV (e.g. Intervet Progard Puppy DPV)
14 weeks	Same as above
16 -18 weeks (<u>optional</u>)	Same as above (<u>optional</u>)
20 weeks or older, if allowable by law	Rabies
1 year	Distemper + Parvovirus, MLV
1 year	Rabies, killed 3-year product (give 3-4 weeks <i>apart</i> from distemper/parvovirus booster)

Perform vaccine antibody titers for distemper and parvovirus every three years thereafter, or more often, if desired. Vaccinate for rabies virus according to the law, except where circumstances indicate that a written waiver needs to be obtained from the primary care veterinarian. In that case, a rabies antibody titer can also be performed to accompany the waiver request. [See www.rabieschallengefund.org](http://www.rabieschallengefund.org)

W. Jean Dodds, DVM